

On Semisimple Hopf Algebras of Dimension pq

Shlomo Gelaki ¹

Department of Mathematics and Computer Science
Ben Gurion University of the Negev
Beer Sheva, Israel

Sara Westreich ¹

Interdisciplinary Department of the Social Science
Bar-Ilan University
Ramat-Gan, Israel

July 23, 1997

In this paper we consider semisimple Hopf algebras of dimension pq over an algebraically closed field k of characteristic 0, where p and q are distinct prime numbers. Masuoka has proved that a semisimple Hopf algebra of dimension $2p$ over k , where p is an odd prime, is either a group algebra or a dual of a group algebra [Ma1]. The authors have pushed the analysis further and obtained the same result for semisimple A of dimension $3p$, where p is a prime greater than 3 [GW]. Thus, a natural conjecture is:

Conjecture 1: Any semisimple Hopf algebra of dimension pq over k , where p and q are distinct prime numbers, is either a group algebra or a dual of a group algebra.

A well known property of A , a finite dimensional semisimple group algebra or a dual of a group algebra, is that it is of *Frobenius type*; that is, the dimension of any irreducible representation of A divides the dimension of A (the definition is due to Montgomery [Mo2]). A special case of one of Kaplansky's conjectures [Kap] is:

Conjecture 2: Any semisimple Hopf algebra of dimension pq over k , where p and q are distinct prime numbers, is of Frobenius type.

In this paper we prove among the rest that Conjecture 1 is equivalent to Conjecture 2 (**Theorem 2.7**).

A major role in the analysis is played by $G(A)$, the group of grouplike elements of A . By [NZ], $|G(A)|$ is either 1, p , q or pq . We prove in **Theorem 2.1** that if $p < q$ then $|G(A)| \neq q$, and if $|G(A)| = p$ then $q = 1(\text{mod } p)$. Consequently, we prove in **Theorem 2.2** that if $|G(A)| \neq 1$ and $q \neq 1(\text{mod } p)$ then A is a group algebra.

¹This research was supported by THE ISRAEL SCIENCE FOUNDATION founded by the Israel Academy of Sciences and Humanities.

Thus Theorem 2.2 suggests a question: When is $|G(A)| \neq 1$? In **Proposition 2.3** we prove that this is guaranteed when A^* is of Frobenius type, and in **Theorem 2.4** we prove that if moreover $q \neq 1 \pmod{p}$ then A is a group algebra. In **Theorem 2.6** we prove that if $|G(A^*)| \neq 1$ and A^* is of Frobenius type, then A is either a group algebra or the dual of a group algebra, and $|G(A)| = p < q$ or pq . The equivalence of Conjectures 1 and 2 is thus a consequence of Proposition 2.3 and Theorem 2.6.

We conclude by using Theorem 2.2 to prove in **Theorem 2.9** that if A is a semisimple Hopf algebra of dimension $5p$, p an odd prime, and if $p = 2$ or $4 \pmod{5}$ or $p \in \{13, 23\}$, then A is a group algebra. Moreover, we obtain in **Theorem 2.10** the same result for semisimple A of dimension $7p$, p a prime, and $p = 6 \pmod{7}$ or $p \in \{17, 31\}$.

1 Preliminaries

In this paper k will always denote an algebraically closed field of characteristic 0.

Recall that a finite dimensional Hopf algebra over k is semisimple if and only if it is cosemisimple [LR].

Let A be a finite dimensional semisimple Hopf algebra over k , and let $\rho_V : A \rightarrow \text{End}_k(V)$ be a finite dimensional representation of A . The afforded character χ_V is given by $\chi_V(a) = \text{tr}(\rho_V(a))$ for all $a \in A$. A character χ_V is called irreducible if the representation V is irreducible. Let $C(A)$ denote the character ring of A ; that is, the k -subalgebra of A^* generated by the characters χ_V of finite dimensional A -modules V . The set of all the irreducible characters forms a basis of $C(A)$ [La]. Zhu has proved that $C(A)$ is semisimple and if $e_{A^*}, e_1, \dots, e_k$ are the primitive idempotents of $C(A)$, where e_{A^*} is an integral of A^* , then

$$\dim A = 1 + \sum_{i=1}^k \dim(e_i A^*) \quad (1)$$

and the dimension of each $e_i A^*$ divides the dimension of A [Z]. Note that $\dim C(A) \geq k + 1$, and equality holds if and only if $C(A)$ is commutative.

Let $f : A \rightarrow A^*$ be the map given by $f(a) = a \rightharpoonup \lambda = \sum \langle a, \lambda_{(2)} \rangle \lambda_{(1)}$ for all $a \in A$, where λ is a non-zero integral of A^* . Recall that f gives a linear isomorphism between $kG(A)$ and the sum of the 1-dimensional ideals of A^* (where $G(A)$ denotes the group of grouplike elements of A), and a linear isomorphism between the center $Z(A)$ of A and $C(A)$. Therefore, using the notation of (1), $\dim(e_i A^*) = 1$ for some i if and only if $G(A) \cap Z(A) \neq \{1\}$.

Let A be a finite dimensional semisimple Hopf algebra over k . Any simple subcoalgebra C_l of A^* has a basis $\{x_{ij}^l | 1 \leq i, j \leq n_l\}$, where $\Delta(x_{ij}^l) = \sum_{k=1}^{n_l} x_{ik}^l \otimes x_{kj}^l$ and $\varepsilon(x_{ij}^l) = \delta_{i,j}$. Note that $n_l = 1$ if and only if $C_l = \{g\}$ for some $g \in G(A)$. Nichols and Richmond have proved that if $\dim A$ is odd then A does not have a 2-dimensional irreducible module [NR], hence

$$\dim A = |G(A)| + \sum_l n_l^2, \quad n_l \geq 3. \quad (2)$$

Now, L is an irreducible left coideal of C_l if and only if

$$L = L_j^l = sp\{x_{kj}^l | 1 \leq k \leq n_l\} \quad (3)$$

for some $1 \leq j \leq n_l$. Similarly, R is an irreducible right coideal of C_l if and only if

$$R = R_k^l = sp\{x_{kj}^l | 1 \leq j \leq n_l\} \quad (4)$$

for some $1 \leq k \leq n_l$. Note that

$$\dim(L_j^l \cap R_k^l) = 1 \quad (5)$$

for any $1 \leq j, k \leq n_l$.

The following theorem will be very useful in the sequel.

Theorem 1.1 [R] *If $H \xrightarrow{i} A \xrightarrow{\pi} H$ is a sequence of finite dimensional Hopf algebra maps where i is injective, π is surjective and $\pi \circ i = id_H$ then there exists $B \subset A$ so that:*

- (i) B is a left H -module algebra and coalgebra via the adjoint action.
- (ii) B is a left H -comodule algebra and coalgebra via $\rho(b) = \sum b^{(1)} \otimes b^{(2)} = \sum \pi(b_{(1)}) \otimes b_{(2)}$.
- (iii) $B \cong A/AH^+$ as a coalgebra, via the map $b \times h \mapsto b\varepsilon(h)$.
- (iv) B is a subalgebra coideal of A .
- (v) As an algebra $A = B \times H$ is a smash product.
- (vi) As a coalgebra $A = B \times H$ is a smash coproduct, that is: $\Delta(b \times h) = \sum b_{(1)} \times b_{(2)}^{(1)} h_{(1)} \otimes b_{(2)}^{(2)} \times h_{(2)}$.
- (vii) The map $B \times H \rightarrow A$ ($b \times h \mapsto bi(h)$) is an isomorphism of bialgebras.

2 Main Results

We start by determining the order of $G(A)$.

Theorem 2.1 *Let A be a semisimple Hopf algebra of dimension pq , where $p < q$ are two prime numbers. Then:*

1. $|G(A)| \neq q$.
2. If $|G(A)| = p$ then $q = 1(\text{mod } p)$.

Proof: 1. Suppose on the contrary that $|G(A)| = q$. If $G(A) \cap Z(A) = G(A)$ then $H = kG(A)$ is central in A , hence is a normal sub Hopf algebra of A . Since $A/(AH^+)$ is a Hopf algebra of dimension p it follows by [Z] that $A/(AH^+) \cong kC_p$, and hence $A \cong kC_q \#_{\sigma} kC_p$. Since kC_q is central the afforded action is trivial [Ma3, Section 1]. Therefore A is isomorphic as an algebra to the twisted group ring $kC_q^t[C_p]$ of the cyclic group C_p over the commutative algebra kC_q . Hence A must be commutative (See the proof of [Ma2, Proposition 2.3(2)]). Thus, A^* is a group algebra and hence of Frobenius type. By (2), $pq = q + ap^2 + bq^2$ for some integers $a, b \geq 0$. But $p < q$, hence $b = 0$, which yields a contradiction. We conclude that $G(A) \cap Z(A) = \{1\}$. Therefore, using the notation of (1), $\dim(e_i A^*) \in \{p, q\}$ for all i . Let E_0 be the integral of $kG(A)$ with $\varepsilon(E_0) = 1$. Since $\dim(A/AH^+) = p$, it follows that $AH^+ = A(1 - E_0)$ has dimension $(q - 1)p$ and thus $\dim(AE_0) = p$. Moreover, $E_0 e_A = e_A$, hence $E_0 = e_A + \sum_j e_{i_j}$. But $p < q$, hence counting dimensions yields a contradiction and the result follows.

2. If $G(A) \cap Z(A) = G(A)$ then $H = kG(A)$ is central in A . As in the proof of part 1, it follows that $A \cong kC_p \#_{\sigma} kC_q$, and hence that A is commutative. Therefore, A^* is a group algebra and hence of Frobenius type. By (2), $pq = p + ap^2 + bq^2$ for some integers $a, b \geq 0$. Clearly, $b = 0$ and hence $q = 1 + ap$. If $G(A) \cap Z(A) = \{1\}$ then, using the notation of (1), $\dim(e_i A^*) \in \{p, q\}$ for all i . Since $\dim(A/AH^+) = q$, it follows that $AH^+ = A(1 - E_0)$ has dimension $(p - 1)q$ and thus $\dim(AE_0) = q$. Hence, $E_0 = e_A + \sum_j e_{i_j}$. But, counting dimensions yields that $\dim(e_{i_j} A^*) = p$ for all j , and the result follows in this case as well. ■

The following is a direct consequence of Theorem 2.1:

Theorem 2.2 *Let A be a semisimple Hopf algebra of dimension pq , where $p < q$ are two prime numbers satisfying $q \not\equiv 1 \pmod{p}$. If $|G(A)| \neq 1$ then A is a group algebra.*

In what follows we find out when $|G(A)| \neq 1$ is guaranteed.

Proposition 2.3 *Let A be a semisimple Hopf algebra of dimension pq , where $p < q$ are two prime numbers. If A^* is of Frobenius type then either $|G(A)| = p$ and $q \equiv 1 \pmod{p}$, or $|G(A)| = pq$.*

Proof: If A is cocommutative then $|G(A)| = pq$. Otherwise, $|G(A)| \neq pq$, and by Theorem 2.1, $|G(A)| \neq q$. If $|G(A)| = 1$, then by (1), $pq = 1 + ap^2 + bq^2$ for some integers $a, b \geq 0$, as A^* is of Frobenius type. But, $q^2 > pq$ hence $b = 0$ which yields a contradiction. ■

As a corollary we have the following:

Theorem 2.4 *Let A be a semisimple Hopf algebra of dimension pq , where $p < q$ are two prime numbers satisfying $q \not\equiv 1 \pmod{p}$. If A^* is of Frobenius type then A is a group algebra.*

We consider now semisimple Hopf algebras A of dimension pq such that A^* is of Frobenius type.

Proposition 2.5 *Let A be a non-cocommutative and non-commutative semi-simple Hopf algebra of dimension pq over k , where $p < q$ are prime numbers. Let $C(A^*)$ be the character ring of A^* . If A^* is of Frobenius type then:*

1. $C(A^*)$ is commutative.
2. As a coalgebra $A = k1 \oplus kg \oplus \cdots \oplus kg^{p-1} \oplus C_1 \oplus \cdots \oplus C_a$, where $a = \frac{q-1}{p}$ and C_i is a simple subcoalgebra of A of dimension p^2 for all $1 \leq i \leq a$.
3. $gC_i = C_i = C_i g$ for all $1 \leq i \leq a$.

Proof: Set $H = kG(A)$. By Theorem 2.1 and Proposition 2.3, $\dim H = p$. If H is central in A then H is a normal sub Hopf algebra of A . Since A/AH^+ is a Hopf algebra of dimension q it follows by [Z] that $A/AH^+ \cong kC_q$, and hence $A \cong kC_p \#_{\sigma} kC_q$. Since kC_p is central the afforded action is trivial [Ma3, Section 1]. Therefore A is isomorphic as an algebra to the twisted group ring $kC_p^t[C_q]$ of the cyclic group C_q over the commutative algebra kC_p . Hence A must be commutative (See the proof of [Ma2, Proposition 2.3(2)]). We conclude that $G(A) \cap Z(A) = \{1\}$.

Set $n = \dim C(A^*) - 1$. Then, by (1) there exist two natural numbers a and b such that:

$$pq = 1 + ap + bq \quad (6)$$

and

$$n \geq a + b. \quad (7)$$

Clearly, $a \geq 1$ and hence $b < q$. Moreover, A^* is of Frobenius type and $p < q$, hence by (2)

$$pq = p^2(n + 1 - p) + p. \quad (8)$$

Substituting (6) and (7) in (8) yields

$$pq \geq p^2(a + b + 1 - p) + p = p^2 \left(\frac{(p-b)q-1}{p} + b + 1 - p \right) + p$$

and hence $(1-p+b)q \geq (1-p+b)p$. Since $p < q$ and $b < p$, this is possible if and only if $b = p-1$ and equality holds in (7). This implies that $C(A^*)$ is commutative and that $pq = 1 + ap + (p-1)q$ and hence $a = \frac{q-1}{p}$. Let

$$e_A, e_1, \dots, e_a, e_{a+1}, \dots, e_{a+p-1}$$

be the primitive idempotents of $C(A^*)$, where e_A is the integral of A with $\varepsilon(e_A) = 1$, $\dim(Ae_i) = p$ for $1 \leq i \leq a$ and $\dim(Ae_{a+j}) = q$ for $1 \leq j \leq p-1$. Let $E_0 = \frac{1}{p} \sum_{i=0}^{p-1} g^i$ be an integral of H . Since $\dim(A/AH^+) = q$, it follows that $AH^+ = A(1 - E_0)$ has dimension $(p-1)q$ and thus $\dim(AE_0) = q$. Moreover, $E_0 e_A = e_A$, hence counting dimensions yields that

$$E_0 = e_A + e_1 + \cdots + e_a.$$

Now, $C(A^*)$ is commutative, thus $\dim(C(A^*)e_A) = \dim(C(A^*)e_i) = 1$ for all $1 \leq i \leq a$ and hence

$$\dim(C(A^*)E_0) = a + 1. \quad (9)$$

Since the set of all the irreducible left A^* -modules consists of p 1-dimensional modules and $a = \frac{q-1}{p}$ p -dimensional modules it follows that

$$A = k1 \oplus kg \oplus \cdots \oplus kg^{p-1} \oplus C_1 \oplus \cdots \oplus C_a$$

as a coalgebra where C_i is a simple subcoalgebra of A of dimension p^2 for all $1 \leq i \leq a$. Let

$$\{1, g, \dots, g^{p-1}, \chi_1, \dots, \chi_a\}$$

be the set of irreducible characters of A^* , where χ_i corresponds to C_i . This set clearly forms a basis of $C(A^*)$. Then

$$C(A^*)E_0 = sp\{E_0, E_0\chi_1, \dots, E_0\chi_a\}$$

which implies by (9) that $\{E_0, E_0\chi_1, \dots, E_0\chi_a\}$ forms a basis of $C(A^*)E_0$. If $g\chi_i = \chi_j$ for $i \neq j$ then $E_0\chi_i = E_0\chi_j$ which is a contradiction. Therefore, $g\chi_i = \chi_i$, and hence

$$gC_i = C_i = C_i g$$

for all i . This concludes the proof of the proposition. ■

Theorem 2.6 *Let A be a semisimple Hopf algebra of dimension pq over k , where $p < q$ are prime numbers. If A^* is of Frobenius type and $|G(A^*)| \neq 1$, then A is either a group algebra or a dual of a group algebra, and $|G(A)| = p < q$ or pq .*

Proof: If A is either cocommutative or commutative then A is either a group algebra or a dual of a group algebra respectively. In any event A^* is of Frobenius type, hence by Proposition 2.3, $|G(A)| = p < q$ or pq and we are done.

Suppose that A is not cocommutative and not commutative. Then Proposition 2.5 is applicable. Set $H = kG(A)$. By Proposition 2.3, $|G(A)| = p$. By Theorem 2.1, $|G(A^*)| \neq q$, hence $|G(A^*)| = p$ too. Thus we have the following sequence of maps:

$$H \xhookrightarrow{i} A \xrightarrow{\pi} H$$

where i is the inclusion map and π is a surjection homomorphism of Hopf algebras. If $\pi \circ i = \varepsilon$ then $H \subseteq K = A^{coH}$. Since K is a left coideal of A , it is a direct sum of irreducible left coideals $K = k1 \oplus kg \oplus \cdots \oplus kg^{p-1} \oplus V_1 \oplus \cdots \oplus V_n$, where $\dim V_i = p$ for all i as A^* is of Frobenius type. This is a contradiction since p does not divide $\dim K = q$. Therefore $\pi \circ i \neq \varepsilon$ and we may assume that $\pi \circ i = id_H$. Therefore by Theorem 1.1, there exists $B \subset A$ so that $A \cong B \times H$. By Theorem 1.1(iv), B is a left coideal of A , hence a direct sum of irreducible left coideals of A . By Proposition 2.5(2), the dimensions of these irreducible left coideals are either 1 or p . Since $\dim B = q$, it follows that B contains an irreducible left coideal V of A , of dimension p . Since $V \subset C$ for some p^2 -dimensional simple subcoalgebra C , it follows by

Theorem 1.1(vii) and Proposition 2.5(3) that $V \times H \subseteq C$. But, $\dim(V \times H) = p^2 = \dim C$, hence $V \times H = C$. By Theorem 1.1(iii), $A/AH^+ \cong B$ as coalgebras, and V is the image of $C = V \times H$ under this isomorphism, hence

V is a subcoalgebra of B .

We wish to prove that V is a simple sub coalgebra of B and thus to reach a contradiction. Note that since V is an irreducible left coideal of A it follows that $V \times g^i$ is also an irreducible left coideal of A for all $0 \leq i \leq p-1$. By Proposition 2.5(3), it follows that the set

$$\{V \times g^i | 0 \leq i \leq p-1\}$$

is the set of all the irreducible left coideals of A contained in C . Since, V is a left coideal of A , it follows from Theorem 1.1(ii) that V is an H sub-comodule of B . Let $\rho : B \rightarrow H \otimes B$ be the comodule structure map, and write $V = \bigoplus_{i=0}^{p-1} V_i$, where $V_i = \rho^{-1}(g^i \otimes V)$. We claim that $\dim V_i = 1$ for all i . Indeed, let $\{v_0, \dots, v_{p-1}\}$ be a basis of V consisting of homogenous elements; that is, $\rho(v_i) = g^{m_i} \otimes v_i$ for some $0 \leq m_i \leq p-1$. Let $0 \neq v \in V$ and write $\Delta_B(v) = \sum_{i=0}^{p-1} b_i \otimes v_i$. Then by Theorem 1.1(vi),

$$\Delta_A(v \times 1) = \sum_{i=0}^{p-1} b_i \times g^{m_i} \otimes v_i \times 1.$$

Therefore, using Kaplansky's notation [Kap], $L(v \times 1) = sp\{b_i \times g^{m_i} | 0 \leq i \leq p-1\} \subset C$ is a **right** coideal of A of dimension $\leq p$. Since C is a simple subcoalgebra of A of dimension p^2 , it follows that $\dim(L(v \times 1)) = p$ and $L(v \times 1)$ is irreducible. Therefore by (5), $\dim(L(v \times 1) \cap (V \times g^i)) = 1$ for all i , hence $\{m_i | 0 \leq i \leq p-1\} = \{0, 1, \dots, p-1\}$. Thus V has a basis $\{v_i | 0 \leq i \leq p-1\}$, where $\rho(v_i) = g^i \otimes v_i$. Since V is an H -comodule coalgebra it follows that $\Delta_B(V_i) \subseteq \sum_{j=0}^{p-1} V_j \otimes V_{i-j}$, hence $\Delta_B(v_i) = \sum_{j=0}^{p-1} \alpha_{ij} v_j \otimes v_{i-j}$ for all i , for some $\alpha_{ij} \in k$. Computing $\Delta_A(v_i \times 1)$ yields that $R_i = L(v_i \times 1) = sp\{\alpha_{ij} v_j \times g^{i-j} | 0 \leq j \leq p-1\} \subset C$ is a right coideal of A of dimension $\leq p$, for all i . Hence $\dim R_i = p$ and

$$R_i = sp\{v_j \times g^{i-j} | 0 \leq j \leq p-1\} \quad (10)$$

is irreducible. It is straightforward to verify that $R_i \neq R_t$ for $i \neq t$, and hence the set $\{R_i | 0 \leq i \leq p-1\}$ is the set of **all** the irreducible right coideals of A which are contained in C .

Finally, let $D \subseteq V$ be a subcoalgebra. By Theorem 1.1(vi), $D \times H \subseteq C$ is a **right** coideal of A and hence $D \times H = \bigoplus_i R_i$, where R_i is as in (10). But, the image of $D \times H$ under the map $id \otimes \varepsilon : A \rightarrow B$ equals D , while the image of $\bigoplus_i R_i$ under this map equals V . Therefore $D = V$, and hence V is a simple coalgebra. But, this is a contradiction since $\dim V = p$ is not a square. ■

As a corollary we obtain the following:

Theorem 2.7 *If both A and A^* are of Frobenius type then A is either a group algebra or the dual of a group algebra.*

Proof: Follows from Proposition 2.3 and Theorem 2.6. ■

We conclude the paper by considering semisimple Hopf algebras of dimensions $5p$ and $7p$.

Lemma 2.8 *Let A be a semisimple Hopf algebra of odd dimension over k . If $|G(A)| = 1$ then there exists an irreducible A^* -module V with $\dim V \geq 4$.*

Proof: Suppose on the contrary that for any non-trivial A^* -irreducible module V , $\dim V \leq 3$. Then by [NR], $\dim V = 3$. Hence, $\dim(V \otimes V^*) = 9$ and by [La], $V \otimes V^* = k \oplus V_1 \oplus \cdots \oplus V_i$, where $V_j \neq k$ is an A^* -irreducible module for all j . Since $\dim V_j = 3$, this is a contradiction. ■

Theorem 2.9 *Let A be a semisimple Hopf algebra over k . If $\dim A = 5p$, p an odd prime, and if $p = 2$ or $4 \pmod{5}$ or $p \in \{13, 23\}$, then A is a group algebra.*

Proof: We wish to show that $|G(A)| \neq 1$. Suppose on the contrary that $|G(A)| = 1$. Set $n = \dim C(A^*) - 1$. By (1), there exist two natural numbers $1 \leq a$ and $1 \leq b \leq 4$ so that

$$\begin{aligned} 5p &= 1 + 5a + bp, \\ n &\geq a + b \quad \text{and} \\ 5p &\geq 9(n - 1) + 16 + 1 \end{aligned}$$

where the last inequality follows from (2) and Lemma 2.8. Hence $(-20 + 9b)p \geq 45b + 31$. But, if $p = 2$ or $4 \pmod{5}$ then $b = 2$ or 1 respectively and if $p \in \{13, 23\}$ then $b = 3$. In any event this is impossible and we have proved that $|G(A)| \neq 1$. The result follows now from Theorem 2.2. ■

Theorem 2.10 *Let A be a semisimple Hopf algebra over k . If $\dim A = 7p$, p a prime, and if $p = 6 \pmod{7}$ or $p \in \{17, 31\}$, then A is a group algebra.*

Proof: Suppose $|G(A)| = 1$ and set $n = \dim C(A^*) - 1$. By (1), there exist two natural numbers $1 \leq a$ and $1 \leq b \leq 6$ so that

$$\begin{aligned} 7p &= 1 + 7a + bp, \\ n &\geq a + b \quad \text{and} \\ 7p &\geq 9(n - 1) + 16 + 1 \end{aligned}$$

where the last inequality follows from (2) and Lemma 2.8. Thus, $(-14 + 9b)p \geq 63b + 47$. But, if $p = 6 \pmod{7}$ or $p \in \{17, 31\}$ then this is impossible. The result follows now from Theorem 2.2. ■

References

- [GW] S. Gelaki and S. Westreich, Semisimple Hopf Algebras of Dimension $3p$, submitted.
- [Kap] I. Kaplansky, Bialgebras, University of Chicago, 1975.
- [La] R. G. Larson, Characters of Hopf Algebras, J. Algebra **17** (1971), 352-368.
- [LR] R. G. Larson and D. E. Radford, Finite Dimensional Cosemisimple Hopf Algebras in Characteristic 0 are Semisimple, J. of Algebra **117** (1988), 267-289.
- [Ma1] A. Masuoka, Semisimple Hopf Algebras of Dimension $2p$, Communication in Algebra **23** No.5 (1995), 1931-1940.
- [Ma2] A. Masuoka, Semisimple Hopf Algebras of Dimension 6, 8, Israel Journal of Mathematics, **92** (1995), 361-373.
- [Ma3] A. Masuoka, Self Dual Hopf Algebras of Dimension p^3 Obtained by Extension, J. of Algebra **178** (1995), 791-806.
- [Mo1] S. Montgomery, Hopf algebras and their actions on rings, CBMS Lecture Notes, Vol. 82, AMS, 1993.
- [Mo2] S. Montgomery, private communication.
- [NR] W. D. Nichols and B. Richmond, The Grothendieck Group of a Hopf Algebra, J. of Pure and Applied Algebra **106** (1996), 297-306.
- [NZ] W. D. Nichols and M. B. Zoeller, A Hopf algebra freeness theorem, Amer. J. of Math. **111** (1989), 381-385.
- [R] D. E. Radford, The Structure of Hopf Algebras with a Projection, J. of Algebra **2** (1985), 322-347.
- [Z] Y. Zhu, Hopf algebras of prime dimension, International Mathematical Research Notices No.1 (1994), 53-59.